

**CLASS: A**

**CLASS B:** Contact medical control for administration of two or more IV sedative medications

**PROTOCOL(S) USED IN: Altered Mental Status, Seizures, Patient Restraint Physical & Chemical**

**PHARMACOLOGY AND ACTIONS:**

- A. Benzodiazepine with antianxiety and sedative effects.
- B. Anticonvulsant

**INDICATIONS:**

- A. Status Epilepticus
- B. Chemical Sedation for transcutaneous pacing
- C. Relief of anxiety

**CONTRAINDICATIONS:**

- A. Hypersensitivity
- B. Acute narrow-angle glaucoma

**SIDE EFFECTS AND NOTES:**

- A. Apnea, N/V, drowsiness, restlessness, confusion, delirium, HTN, hypotension
- B. **Class D pregnancy category**; may cause fetal damage
- C. When administering, dilute medication 1:1 with NS prior to administration.

**ADULT DOSING:**

**Status epilepticus**

**2-4 mg IM or slow IV/IO** over 2 minutes. May repeat once if still seizing after 5-10 minutes. Max of 8 mg. Contact OLMC for further doses PRN.

**Chemical Sedation**

**2 mg IM or 1 mg IV**

**Severe Anxiety**

**1 mg IN, IM or slow IV**

**PEDIATRIC DOSING:**

**Status epilepticus**

Consider **Ativan** (28 days to 12 years) dose: 0.05-0.1mg/kg IV/IO/IM over 2-5 mins. IV diluted 1:1 with Normal Saline

- i. If still seizing after 5-10mins you can repeat dose once