

Endotracheal Intubation RSI – 30.061

OBJECTIVES:

- A. To facilitate orotracheal intubation
- B. To protect from increased ICP associated with direct laryngoscopy.
- C. To reduce the discomfort and trauma of intubation in conscious patients.

INDICATIONS:

Patient meets indications previously noted in the orotracheal intubation protocol AND:

- A. Clenched jaw or active gag reflex.
- B. Combativeness threatens the airway, spinal cord stability, and/or transport safety.
- C. The patient is conscious.

CONTRAINDICATIONS:

- A. Inability to ventilate adequately with a bag-valve mask in the event of failed intubation.

PROCEDURE:

- A. Prepare, position, and pre-oxygenate as outlined in the orotracheal intubation protocol.
- B. Induction agents. *Give only one.*
 - a. **Etomidate 0.3 mg/kg IV/IO** push. Single max dose of 30 mg.
 - b. **Ketamine 1-2 mg/kg IV** push. Single max dose of 200 mg.
 - c. **Midazolam 0.1 mg/kg IV/IO** push. Single max dose of 10 mg.
- C. Paralytic agents. *Give only one.*
 - a. **Succinylcholine 1.5 mg/kg IV/IO**. See contraindications below.
 - b. **Rocuronium 1 mg/kg IV/IO**.
 - c. **Vecuronium 0.1 mg/kg IV/IO**.
- D. Adjuncts
 - a. NO DESAT: Increase nasal cannula oxygen to 15 LPM AFTER medications are given.
- E. Assess for apnea and jaw relaxation and gently intubate in a controlled but timely manner when patient becomes relaxed.
- F. Confirm ETT placement, reassess vitals and document as outlined in the orotracheal protocol.
- G. Continued sedation and analgesia are paramount. Continue paralysis as needed.
 - a. **Midazolam 0.05-0.1 mg/kg IV**. Single max dose of 5 mg.
 - b. **Ketamine 0.5-1 mg/kg IV**.
 - c. **Fentanyl 1-2 mcg/kg IV**.
 - d. **Rocuronium 0.3-0.5 mg/kg IV**.
 - e. **Vecuronium 0.1 mg/kg IV**.

SUCCINYLCHOLINE CONTRAINDICATIONS

- A. Crush or burn injuries more than 24 hours old (due to potential for hyperkalemia).
- B. Penetrating eye injuries (relative) due to increased intraocular pressure.
- C. Medical history including malignant hyperthermia, myasthenia gravis, muscular dystrophy, dialysis patient if potassium level is not known, or hyperkalemia.
- D. Hypersensitivity to the drug.

COMMENTS

- A. Repeat boluses of Etomidate should NOT be used for maintenance of sedation after intubation secondary to potential adrenal suppression.
- B. Consider sedation utilizing Ketamine for those patients in whom difficult airway is suspected or those patients with suspected lower airway obstruction: i.e. status asthmaticus, COPD, or severe bronchiolitis.

COMPLICATIONS

- A. Cardiac dysrhythmias.
- B. Hyperkalemia.
- C. Fasciculation's from paralysis.
- D. Vomiting and/or aspiration.
- E. Esophageal intubation – unrecognized esophageal intubation is a “never event”.
- F. Prolonged paralysis & malignant hyperthermia.
- G. Oral trauma.

DOCUMENTATION

- A. As per Orotracheal Intubation protocol.
- B. RSI and sedation/analgesia medications given

PEDIATRIC Rapid Sequence Intubation (RSI)

Procedure:

- A. Prepare, position and pre-oxygenate
- B. Adjuncts
 - a. NO DESAT: increase NC oxygen to 15 lpm AFTER medications are given
 - b. RSI for pediatrics 2 years and under, consider **Atropine 0.02 mg/kg IV/IO**
- C. Induction agent *Give only one*
 - a. **Ketamine – 1 mg/kg IV/IO**
 - b. **Etomidate – 0.3 mg/kg IV/IO**
- D. Paralytic agent *Give only one*
 - a. **Succinylcholine – 1.5-2 mg/kg IV/IO (depending on age)**
 - b. **Rocuronium – 1 mg/kg IV/IO**
 - c. **Vecuronium – 0.1 mg/kg IV/IO**
- E. Assess for apnea and jaw relaxation and gently intubate in a timely manner
- F. Confirm ETT placement, reassess vitals and document as outlined in the orotracheal intubation protocol.
- G. Continued sedation and analgesia are paramount. Continue paralysis PRN
 - a. **Midazolam – 0.1 mg/kg IV/IO and/or**
 - b. **Fentanyl – 0.5 mcg/kg IV/IO**