


# Neonatal Resuscitation – 10.160



## Neonatal Resuscitation – 10.160

### NOTES & PRECAUTIONS:

- A. Neonatal age is birth to 28 days.
- B. Do not use atropine in neonatal resuscitation.
- C. If meconium is lightly stained and infant is vigorous (strong respiratory effort, good muscle tone, heart rate > 100 bpm) endotracheal suctioning should not be performed.
- D. An infant may need resuscitation if intrapartum risk factors for asphyxia are present (prolapsed cord, painful bleeding, prolonged rupture of membranes, maternal fever, multiple births, abnormal presentation, maternal hypotension or seizure)

### KEY CONSIDERATIONS:

Fetal presentation, recent trauma, maternal health/risk factors, maternal medications, previous birth difficulties, APGAR

APGAR SCORE:	0	1	2
Appearance	Blue/Pale	Body pink, blue extremities	Completely pink
Pulse	Absent	Slow (< 100 bpm)	≥ 100 bpm
Grimace	No response	Grimace	Cough or sneeze
Activity	Limp	Some flexion	Active motion
Respirations	Absent	Slow, irregular	Good, crying

### Newborn Target Spo2 after birth:

<b>1 min</b>	<b>60-65%</b>
<b>2 min</b>	<b>65-70%</b>
<b>3 min</b>	<b>70-75%</b>
<b>4 min</b>	<b>75-80%</b>
<b>5 min</b>	<b>80-85%</b>
<b>10 min</b>	<b>85-95%</b>